

FDA Approves Merck's CMV Drug Prevyimis (Letermovir)

The drug is approved to cut the risk of CMV reactivation following a stem cell transplant.

December 14, 2017

The Food and Drug Administration (FDA) has approved Merck's cytomegalovirus (CMV) treatment Prevyimis (letermovir) for adults infected with the virus for use during the period following a stem cell transplant. The drug can be given as once-daily tablets or as an intravenous infusion.

People who test positive for CMV and who receive a stem cell transplant and experience associated suppressed immune system following the procedure are at a high risk for the virus to reactivate and have a higher risk of death as a result of their infection. CMV is common among people living with HIV and was a major cause of death for the population before the advent of effective antiretroviral treatment for the latter virus.

The recommended dose of Prevyimis is 480 milligrams once daily, started at any point between the day of a stem cell transplant and four weeks post-transplant and continued through day 100 after the transplant.

The FDA approval was based on findings from a multicenter, double-blind placebo-controlled Phase III trial of 565 CMV-positive adults who received a stem cell transplant. They were randomized two-to-one to receive either 480 mg of Prevyimis daily (or 240 mg daily if they also received the immunosuppressant drug cyclosporine) or a placebo. The participants were started on treatment within the first four weeks following their transplant and kept on treatment through week 14 post-transplant. They were followed for 48 weeks total.

By week 24 after the transplant, among those who received Prevyimis and those who received the placebo, a respective 18 percent and 42 percent developed a clinically significant CMV infection—defined as CMV end-organ disease or the initiation of preemptive CMV treatment with antivirals, treatment that was prompted by a documented CMV viral load and the participant's clinical condition.

At week 24 post-transplant, 12 percent of those who received Prevyimis had died of any cause, compared with 17 percent of those who received the placebo. At week 48 post-transplant, the corresponding respective figures were 24 percent and 28 percent.

Thirteen percent of those who received Prevymis experienced an adverse cardiac health event compared with 6 percent of those who received the placebo. The most common such events, which were mild or moderate in severity, were tachycardia (rapid heart rate), reported in a respective 4 percent and 2 percent of participants, and atrial fibrillation (rapid and often irregular heartbeat, often including palpitations), reported in a respective 3 percent and 1 percent of participants.

The rate of adverse health events reported by at least 10 percent of those receiving Prevymis and that occurred at a frequency at least 2 percentage points greater than those receiving the placebo were nausea (27 percent among those receiving Prevymis versus 23 percent among those receiving the placebo), diarrhea (26 percent versus 24 percent), vomiting (19 percent versus 14 percent), peripheral edema (14 percent versus 9 percent), cough (14 percent versus 10 percent), headache (14 percent versus 9 percent), fatigue (13 percent versus 11 percent) and abdominal pain (12 percent versus 9 percent).

To read a press release about the FDA approval, [click here](#).