FDA Approves Axicabtagene Ciloleucel for Large B-Cell Lymphoma

Yescarta is the first chimeric antigen receptor T cell (CAR-T) immunotherapy approved for adults.

October 25, 2017 By Food and Drug Administration (FDA)

On October 18, 2017, the Food and Drug Administration granted regular approval to axicabtagene ciloleucel (YESCARTA, Kite Pharma, Inc.) for the treatment of adult patients with relapsed or refractory large B-cell lymphoma after two or more lines of systemic therapy, including diffuse large B-cell lymphoma (DLBCL) not otherwise specified, primary mediastinal large B-cell lymphoma, high-grade B-cell lymphoma, and DLBCL arising from follicular lymphoma.

Axicabtagene ciloleucel is a chimeric antigen receptor (CAR) T cell immunotherapy. It consists of autologous T cells that are genetically modified to produce a CAR protein, allowing the T cells to identify and eliminate CD19-expressing normal and malignant cells.

Approval was based on a single-arm multicenter trial of 108 adult patients with aggressive B-cell non-Hodgkin lymphoma. Eligible patients had refractory disease to the most recent therapy or relapse within one year after autologous hematopoietic stem cell transplantation. Patients received a single infusion of axicabtagene ciloleucel following completion of lymphodepleting chemotherapy.

Of the 101 patients evaluated for efficacy, the objective response rate (ORR) as assessed by independent central review was 72%, with a complete remission (CR) rate of 51% (95% CI: 41, 62). The duration of response (DOR) was longer in patients with a best overall response of CR, as compared to a best overall response of partial remission (PR). Among patients achieving CR, the estimated median DOR was not reached (95% CI: 8.1 months, not estimable [NE]) after a median follow-up of 7.9-months. The estimated median DOR among patients in PR was 2.1 months (95% CI: 1.3, 5.3).

The most common grade 3 or higher adverse reactions (incidence of 10% or greater) include febrile neutropenia, fever, cytokine release syndrome (CRS), encephalopathy, infections, hypotension, and hypoxia. Serious adverse reactions occurred in 52% of patients and included CRS, neurologic toxicity, prolonged cytopenias (including neutropenia, thrombocytopenia, and anemia), and serious infections. Fatal cases of CRS and neurologic toxicity occurred. FDA approved axicabtagene ciloleucel with a Risk Evaluation and Mitigation Strategy.

The recommended dose of axicabtagene ciloleucel is a single intravenous infusion with a target of

2 x 10^6 CAR-positive viable T cells per kg body weight (maximum 2 x 10^8), preceded by fludarabine and cyclophosphamide lymphodepleting chemotherapy. Axicabtagene ciloleucel is not indicated for the treatment of patients with primary central nervous system lymphoma.

Full prescribing information is available at: https://www.fda.gov/BiologicsBloodVaccines/CellularGeneTherapyProducts/ApprovedProducts/ucm581222.htm.

FDA granted orphan drug designation and priority review to axicabtagene ciloleucel for this indication. Approval was granted approximately 6 weeks prior to the due date. A description of FDA expedited programs is in the Guidance for Industry: Expedited Programs for Serious Conditions-Drugs and Biologics, available at: http://www.fda.gov/downloads/drugs/guidancecomplianceregulatoryinformation/guidances/ucm358301.pdf.

Healthcare professionals should report all serious adverse events suspected to be associated with the use of any medicine and device to FDA's MedWatch Reporting System by completing a form online at http://www.fda.gov/medwatch/report.htm, by faxing (1-800-FDA-0178) or mailing the postage-paid address form provided online, or by telephone (1-800-FDA-1088).

This announcement was originally published on the Food and Drug Administration website on October 25, 2017.

© 2026 Smart + Strong All Rights Reserved.

http://beta.docker.cancerhealth.com/blog/fda-approves-axicabtagene-ciloleucel-large-bcell-lymphoma