

FDA Extends Approval of Keytruda for Classical Hodgkin Lymphoma

The checkpoint inhibitor is approved for adults and children with relapsed lymphoma.

October 16, 2020 By [Food and Drug Administration \(FDA\)](#)

On October 14, 2020, the Food and Drug Administration extended the approval of pembrolizumab (KEYTRUDA, Merck Sharp & Dohme Corp.) for the following indications:

- adult patients with relapsed or refractory classical Hodgkin lymphoma (cHL) and
- pediatric patients with refractory cHL, or cHL that has relapsed after 2 or more lines of therapy.

Approval was based on KEYNOTE-204 (NCT02684292), a phase 3, randomized, open-label trial in 304 adult patients with relapsed or refractory cHL after at least one multiagent regimen. Patients were randomized (1:1) to receive either pembrolizumab 200 mg every 3 weeks or brentuximab vedotin (BV) 1.8 mg/kg every 3 weeks for up to 2 years.

Efficacy was based on progression-free survival (PFS) per blinded independent central review assessment. PFS was statistically significantly longer in the pembrolizumab arm. The median PFS was 13.2 months (95% CI: 10.9, 19.4) in the pembrolizumab arm and 8.3 months (95% CI: 5.7, 8.8) in the BV arm, with a hazard ratio of 0.65 (95% CI: 0.48, 0.88; $p=0.0027$).

Serious adverse reactions occurred in 30% of the patients who received pembrolizumab. Serious adverse reactions in $\geq 1\%$ of patients included pneumonitis, pneumonia, pyrexia, myocarditis, acute kidney injury, febrile neutropenia, and sepsis.

Adverse reactions in $\geq 20\%$ of pembrolizumab recipients included upper respiratory tract infection, musculoskeletal pain, diarrhea, cough, pyrexia, fatigue, and rash. Thirty-eight percent of patients had adverse reactions requiring systemic corticosteroids, including pneumonitis in 11%.

The recommended pembrolizumab dose for patients with lymphoma is 200 mg every 3 weeks or 400 mg every 6 weeks intravenously for adults, or 2 mg/kg (up to 200 mg) every 3 weeks intravenously for pediatric patients, for up to 2 years.

[View full prescribing information for Keytruda.](#)

The FDA collaborated with the Australian Therapeutics Goods Administration (TGA) and Health Canada (HC) as part of [Project Orbis](#). The review of the application is ongoing for the TGA and HC. This review used the [Assessment Aid](#), a voluntary submission from the applicant to facilitate the FDA's assessment.

This application was granted orphan drug designation, breakthrough therapy designation, and priority review. A description of FDA expedited programs is in the [Guidance for Industry: Expedited Programs for Serious Conditions-Drugs and Biologics](#).

Healthcare professionals should report all serious adverse events suspected to be associated with the use of any medicine and device to FDA's [MedWatch Reporting System](#) or by calling 1-800-FDA-1088.

For assistance with single-patient INDs for investigational oncology products, healthcare professionals may contact OCE's [Project Facilitate](#) at 240-402-0004 or email OncProjectFacilitate@fda.hhs.gov.

This announcement was originally published on the Food and Drug Administration website on October 14, 2020.

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.cancerhealth.com/blog/fda-extends-approval-keytruda-classical-hodgkin-lymphoma>