

Many Types of Leisure Time Activity May Lower the Risk of Death for Older Adults

Running was associated with the greatest reduction in risk of cancer deaths, according to a new study. Here are other findings.

September 15, 2022 By [National Cancer Institute](#)

Older adults who participate weekly in many different types of leisure time activity, such as walking for exercise, jogging, swimming laps, or playing tennis, may have a lower risk of death from any cause, as well as death from cardiovascular disease and cancer, [according to a new study](#) led by researchers at the National Cancer Institute, part of the National Institutes of Health.

The findings suggest that it's important for older adults to engage in leisure time activities that they enjoy and can sustain, because many types of these activities may lower the risk of death, the authors wrote.

The findings appeared Aug. 24 in JAMA Network Open.

Using data from 272,550 adults between ages 59 and 82 who had completed questionnaires about their leisure time activities as part of the NIH-AARP Diet and Health Study, the researchers looked at whether participating in equivalent amounts of seven different exercise and recreational activities—including running, cycling, swimming, other aerobic exercise, racquet sports, golf, and walking for exercise—was associated with lowered risk of death.

The researchers found that achieving the recommended amount of physical activity per week through any combination of these activities was associated with a 13% lower risk of death from any cause compared with no participation. When they looked at the role of each activity individually, playing racquet sports was associated with a 16% reduction in risk and running with a 15% reduction. However, all activities were similarly associated with lower risks of death.

The second edition of the Physical Activity Guidelines for Americans recommends that adults engage in 2.5 to 5 hours of moderate-intensity aerobic physical activity, or 1.25 to 2.5 hours of vigorous-intensity aerobic physical activity, each week.

The levels of activity by the most active individuals (those who exceeded the recommended levels of physical activity) were associated with even greater reductions in the risk of death, but there

were diminishing returns as activity levels increased. Even people who did some recreational activity, though less than the recommended amount, had a 5% reduction in risk of death than those who did not participate in any of the activities studied.

These activities were also associated with a lower risk of death from cardiovascular disease and cancer. Playing racquet sports was associated with the greatest reduction in risk of cardiovascular deaths (27% reduction), while running was associated with the greatest reduction in risk of cancer deaths (19% reduction).

[This post](#) was originally published August 24, 2022, by the National Cancer Institute. It is republished with permission.

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.cancerhealth.com/blog/many-types-leisure-time-activity-may-lower-risk-death-cancer-older-adults>