

# Older Biologic Age Linked to Elevated Breast Cancer Risk

National Institutes of Health scientists use epigenetics to help predict disease development.

March 1, 2019 By [National Cancer Institute](#)

---

Originally published by the NIH.

Biologic age, a DNA-based estimate of a person's age, is associated with future development of breast cancer, according to scientists at the National Institutes of Health. Biologic age was determined by measuring DNA methylation, a chemical modification to DNA that is part of the normal aging process. The study showed for every five years a woman's biologic age was older than her chronologic or actual age, known as age acceleration, she had a 15 percent increase in her chance of developing breast cancer. The study was published online February 22 in the Journal of the National Cancer Institute.

Scientists from the National Institute of Environmental Health Sciences (NIEHS), part of NIH, speculate that biologic age may be tied to environmental exposures. If so, it may be a useful indicator of disease risk. They used three different measures, called epigenetic clocks, to estimate biologic age. These clocks measure methylation found at specific locations in DNA. Researchers use these clocks to estimate biologic age, which can then be compared to chronologic age.

The researchers used DNA from blood samples provided by women enrolled in the NIEHS-led [Sister Study](#), a group of more than 50,000 women in the U.S. and Puerto Rico. The study was specifically designed to identify environmental and genetic risk factors for breast cancer. The research team measured methylation in a subset of 2,764 women, all of whom were cancer-free at the time of blood collection.

"We found that if your biologic age is older than your chronologic age, your breast cancer risk is increased. The converse was also true. If your biologic age is younger than your chronologic age, you may have decreased risk of developing breast cancer," said corresponding author Jack Taylor, MD, PhD, head of the NIEHS Molecular and Genetic Epidemiology Group. "However, we don't yet know how exposures and lifestyle factors may affect biologic age or whether this process can be reversed."

Lead author Jacob Kresovich, PhD, a postdoctoral fellow in the Taylor group, had read studies that used epigenetic clocks to predict age-related mortality. Since age is the leading risk factor for breast cancer, he hypothesized that age acceleration may be associated with higher breast cancer

risk.

“If you look at a group of people who are all the same age, some may be perfectly healthy while others are not,” Kresovich said. “That variability in health may be better captured by biologic age than chronologic age.”

Kresovich suggests that using DNA methylation to measure biologic age may help scientists better understand who is at risk for developing cancer and other age-related diseases. This research is an example of epigenetics, a field that studies how biochemical processes turn individual genes on or off, without affecting the DNA sequence.

The Taylor group plans to continue using epigenetic data, along with information on genetics, environment, and lifestyle to better understand how these factors interact and contribute to disease risks.

[This news release](#) was originally published by the National Institutes of Health on February 22, 2019.

---

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.cancerhealth.com/blog/older-biologic-age-linked-elevated-breast-cancer-risk>